



## Update on Fish Disease Situation in Italy

**Vendramin, Niccolò; Toffan, A.**

*Published in:*

16th Annual Meeting of the National Reference Laboratories for Fish Diseases

*Publication date:*

2012

*Document Version*

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Vendramin, N., & Toffan, A. (2012). Update on Fish Disease Situation in Italy. In *16th Annual Meeting of the National Reference Laboratories for Fish Diseases*

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# UPDATE ON FISH DISEASE SITUATION IN ITALY

Vendramin N.<sup>1</sup> & Toffan A.<sup>2</sup>

<sup>1</sup>DTU Vet National Veterinary Institute, Høngøvej 2 8200 Aarhus N, [niven@dtu.vet.dk](mailto:niven@dtu.vet.dk)

<sup>2</sup>IZSVenezie, Viale dell'Università 10 Legnaro Padova, [atoffan@izsvenezie.it](mailto:atoffan@izsvenezie.it)

---

## Abstract:

In this presentation we provide a general overview of the aquatic animal health issues related to the aquaculture sector and wild environment in Italy in 2011.

Considering saltwater species European sea bass (*Dicentrarchus labrax*) and Gilthead sea bream (*Sparus aurata*) are still the most widely farmed species, nevertheless some “new” candidates are employed often.

The first two species represents more than 95% of the total production while the remaining is obtained by different promising candidates species, including sole (*Solea solea*), meagre (*Argyrosomus regius*), northern blue fin tuna (*Thunnus thynnus*) and amberjack (*Seriola dumerilii*) for which some breeding/reproduction plans have been attempted by some hatcheries.

The farming of sea bass and sea bream is affected by the presence of several important diseases.

Firstly considering Bacterial diseases, Marine Flexibacteriosis, caused by *Tenacibaculum maritimum*; Vibriosis, caused by *Listonella anguillarum* and Pasteurellosis caused by *Photobacterium Damselfa subsp. piscicida* are considered the major bacterial diseases even though for these pathogens exists chemicals and for some of them efficacious vaccines.

Considering viral diseases Viral encephalopathy and retinopathy (VER) still plays a key role in some areas where, mortalities ranging from 30-40% can be observed in sea bass rearing unit.

Nevertheless in recent years, serious epizootics affecting sea bream larvae, previously considered a resistant species, have been reported. Finally the appearance of the clinical disease has been recorded also in the wild.

Lymphocystis disease (LCD) represents an important disease not for its pathogenicity but for the interferences with strict production plans of farm.

Finally, considering parasites, over than “old” protozoans (*Cryptocaryon irritans* and *Amyloodinium ocellatum*) and gill flukes (*Diplectanum aequans* and *Sparicotyle chrysophrii*) mainly present in inland farms (earth ponds and concrete tanks based); isopods crustacea (*Ceratotoa*, *Anylocra*) and Enteromyxidiosis (*Enteromyxum leei*) represent a treat for offshore cages.

Among the dismetabolic unknown aetiology disease Winter Syndrome, affects mainly 1-year-old sea bream causing mortalities ranging from 5-15%. The therapeutic treatment addresses great attention to the diet, especially at the end of Summer and the approaching winter season.

Considering trout farming which is a well developed industry two main pathological scenarios are present. Farms with low water temperature (mainly located in the mountains) can be more affected and damaged by viral diseases (i.e. viral haemorrhagic septicaemia VHS) which is one of the most important problem.

Rainbow trout fry syndrome (RTFS) is responsible for significant mortalities in salmonids, during juvenile stages, particularly if not treated promptly.

Another bacterial problem that seems to be re-emerging is Enteric Red mouth (*Yersinia ruckerii*).

Finally considering freshwater wild environment a mortality outbreak in eels is presented.